

AD-A111 186

UNIVERSITY OF SOUTHERN CALIFORNIA LOS ANGELES DEPT O--ETC F/6 28/10
DIAGNOSTICS AND CHEMICAL APPLICATIONS OF MULTIPHOTON ABSORPTION--ETC(U)
SEP 81 S W BENSON APOSR-77-3879

UNCLASSIFIED

APOSR-TR-82-0063

NL

for
AD-111



END
DATE
FILMED
9-82
DTIC

LEVEL

④

FINAL SCIENTIFIC REPORT

AD A111155

"DIAGNOSTICS AND CHEMICAL APPLICATIONS OF
MULTIPHOTON ABSORPTION PROCESSES"

by

Principal Investigator: Sidney W. Benson
Professor of Chemistry
Department of Chemistry
University of Southern California
Los Angeles, California 90007

DTIC
ELECTE
FEB 22 1982
S H

AFOSR-77-3279

Sponsored by:

Air Force Office of Scientific Research
Directorate of Chemical Sciences
AFOSR/NC
Bolling Air Force Base, D.C. 20332

DTIC FILE COPY

1 April 1977 - 30 September 1981

82 0.


I. Objective

To examine and analyze quantitatively the mechanism and parameters of the processes of multiphoton absorption from intense laser fields and to observe and measure the intramolecular dynamic processes following such absorption.

II. Method

It was the original objective of the present program to develop a simple molecular beam system for examining the effect of various laser parameters on the rate of energy absorption by irradiated molecules. In the course of developing this system we discovered the utility of using isotopic branching to accomplish a similar purpose and to that end we have explored the multiphoton dissociation of 2-deuterioethyl chloride, $\text{CH}_2\text{DCH}_2\text{Cl}$ ($\text{C}_2\text{H}_4\text{DCl}$).


Excitation of $\text{C}_2\text{H}_4\text{DCl}$ above the activation threshold of about 56 kcal leads to production of HCl or DCl in a concerted molecular process:



The branching ratio $R = k_1/k_1'$ can be measured from the final products and is a monotonic function of the energy of the decomposing molecules. We have made a careful study [ref.5] of the thermal system from 700 - 1000°K and find the Arrhenius relation:

$$k_1/k_1' = 1.44 (\pm 0.05) \exp \left\{ (1500 (\pm 50 \text{ cal/mol}^\circ\text{K})/RT) \right\}$$

We have also modeled the system (RRKM Theory) and shown that the model fits the absolute rates in the pressure dependent region. This



permits us to assign a mean energy content E to each value of R observed and to fix the mean time of decomposition of the decomposing molecule [Ref. ^{3,5}2,4,6]. In this way the observed branching ratio provides a "chemical clock" for the last excitation and decomposition events. At low pressures (< 100 mtorr, $t_{\text{collision}} \geq 10^{-5.5}$ sec) the average decomposing molecule had an internal energy of 88 ± 6 kcal and decomposed in $10^{-9 \pm 0.5}$ sec. It thus absorbed its last photon in about 1 to 3 nanoseconds.

On studying the effects of pressure [Ref. ^{2,4,5}2,5,6] on this system we discovered the bimodal response to the laser pulse shape. A typical laser pulse has an intense; short peak ($0.1 \mu\text{sec}$) and a long, weaker tail ($\sim 1 \mu\text{sec}$) each with about $1/2$ the total energy. Each pumps at a different rate and produces a different energy population. At 30 torr Ne (or He) we see only the effects of the short peak, the slowly pumped molecules in the tail being quenched by collisions before dissociation. The branching ratio R_p corresponding to this peak have energies of 130 ± 15 kcal/mole and life-times of about $10^{-11 \pm 0.5}$ sec. They absorbed their last photon in 10^{-11} sec and have experienced pumping and decomposition in a single mode pulse during a time of about 5 nanosec. To do this the energized molecule had to have an absorption cross-section about 100 times larger than the cross-section of room temperature molecules.

We feel that the discovery of isotopic branching as a chemical clock provides an important new tool to laser scientists. It makes it possible to place intramolecular laser events on an absolute time scale and to begin to ask and answer questions about the time scale for the flow of energy in molecules, both large (> 6 atoms) and very large.

MAJOR ACHIEVEMENTS

- (A) Establishment of isotope branching as chemical clock and energy measure via RRKM
- (B) Connection of Thermal—Fall-off and RRKM with MPD via RRKM
- (C) Discovery of huge changes in X-section with energy
- (D) Discovery of peak and tail effects
- (E) Operations of molecular beam with pulse inlet and IR and excimer laser irradiation

ACTIVE PERSONNEL

- 1.) Dr. A. J. Colussi
- 2.) Dr. J. C. Diels
- 3.) Dr. P. J. Evans
- 4.) Dr. P. Papagiannakopoulos
- 5.) Professor Choi (Korea)
- 6.) Mr. Ken Kosnik (grad student)

PUBLICATIONS

- 1.) "Molecular Beam Sampling", A. J. Colussi and Sidney W. Benson, Int. J. Chem. Kin. 10, 1091 (1978).
- 2.) "Intramolecular Isotope Effects in Laser Multiphoton Dissociation of $\text{CH}_2\text{DCH}_2\text{Cl}$ ", A. J. Colussi, R. J. Hwang, J. Tise and Sidney W. Benson, Chem. Phys. Letters, 52, 349 (1977).
- 3.) "The Prediction of Thermochemical and Kinetic Data for Gas Phase Reactions". (Plenary Lecture). 4th Int. Symp. on Plasma Chemistry, S. W. Benson, Pure and Applied Chem., 52, 1767 (1980).
- 4.) "Intramolecular Isotopic Effect in the Pyrolysis of Ethyl-2d₁ Chloride, P. J. Papagiannakopoulos and Sidney W. Benson, Int. J. Chem. Kin. (1982).
- 5.) "The Effects of Buffer Gases in the Infrared Multiphoton Dissociation of $\text{C}_2\text{H}_4\text{DCl}$. A Chemical Clock to Explore Highly Excited Molecules", P. J. Papagiannakopoulos, Ken Kosnik and Sidney W. Benson, Int. J. Chem. Kin. (1982).

REPORT DOCUMENTATION PAGE		22-APR-81 READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFOSR-TR-82-0063	2. GOVT ACCESSION NO. AD-A111155	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) DIAGNOSTICS AND CHEMICAL APPLICATIONS OF MULTI-PHOTON ABSORPTION PROCESSES		5. TYPE OF REPORT & PERIOD COVERED FINAL	
		6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR Sidney W. Benson		8. CONTRACT OR GRANT NUMBER AFOSR-77-3279	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Chemistry Department University of Southern California Los Angeles, CA 90007		10. PROGRAM ELEMENT PROJECT TASK AREA & WORK UNIT NUMBERS 61102F 2303/B1	
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Air Force /NC Air Force Office Scientific Research Bolling Air Force Base, D. C. 20332		12. REPORT DATE Sep 81	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 3	
		15. SECURITY CLASS. (of this report) Unclassified	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE			
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. SUBJECT TERMS (Continue on reverse side if necessary and identify by block number) Multiphoton Dissociation - isotopic branching ratio Laser - deuterio ethyl chloride, isotope effects collisional quenching			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Isotopic branching ratios are shown to be a very sensitive tool for the quantitative analysis of the absolute time scale and total energy content of molecules undergoing photolysis in an IR laser beam. A study and RRKM analysis has been done of the thermal decomposition of deuterio-ethyl chloride over the range 700-1100°K and at high and low pressures (fall-off region). This has been applied to analysis of the multiphoton dissociation with a CO₂ laser at varying pressures. It is shown that the laser pulse produces two independent effects due to its long, low intensity tail following the high intensity peak. In the latter			

DD FORM 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Molecules absorb about 85 kcal and decompose in $10^{-8.5}$ sec. Also of importance is the finding that energized molecules can have IR absorption cross-sections an order of magnitude or more greater than their cross-section at room temperature.

10 to $10^{-8.5}$ sec

unclassified

END

DATE
FILMED

3-82

DTIC